

SUPREP Instructions

General Procedure Information

Please read these instructions two weeks before

- Arrange to have a driver to come with you and remain in our waiting room during your test.

Laxative Prep Method

***Follow these instructions. Do NOT follow instructions on the box**

SUPREP – Split dose

Supplies to purchase

1. SUPREP kit (fill prescription)
2. Clear liquid beverages/product

General Instructions

2 weeks before procedure

- Stop taking herbal products
- Stop eating snack containing olestra (found in “light” snacks)

1 week before procedure

- Stop taking aspirin, NSAIDs and blood Thinners 3, 4, or 5 days or as approved by your prescribing physician. You may take Tylenol (acetaminophen)
- Stop multivitamins, vitamin E, iron and fish oil supplements

Days before your procedure

- Purchase supplies
- Read through these directions
- If your procedure is at Summit Endoscopy Center, expect call from anesthesia department to discuss your medical history.

If you are diabetic, read and follow the special diabetic instructions. Questions? Please see frequently asked questions first. If you still have questions, please call.

Day Before Procedure

DIET INSTRUCTIONS

- You may only have CLEAR LIQUIDS the entire day before your procedure
- **NO SOLID FOODS** should be eaten on this day
- Examples of CLEAR LIQUIDS: strained fruit juices (apple, white grape, white cranberry). Limeade, lemonade, coffee, tea (no creamer), Coke, Pepsi, Diet soda, non-red soft drinks, chicken or beef broth, gelatin desserts without added fruit or whipped topping
- AVOID RED and PURPLE liquids, including food dye
- AVOID MILK and ALCOHOLIC beverages

LAXATIVE PREP INSTRUCTIONS

- At 5 PM, Pour one 6 ounce bottle of SUPREP liquid into the mixing container. Add cool drinking water to the 16 ounce line on the container.
- Drink ALL the liquid in the container over a 30 minute period.
- Immediately follow with 32 ounces of clear liquids over the next hour.
- Continue to drink clear liquids until bedtime.

Day of Procedure

LAXATIVE PREP INSTRUCTIONS

6 hours before leaving for the procedure

- Pour one 6 ounce bottle of SUPREP liquid into the mixing container. Add cool drinking water to the 16 ounce line on the container.
- Drink ALL the liquid in the container over a 30 minute period.
- Immediately follow with 32 ounces of clear liquids over the next hour.
- **Put nothing else in your mouth** except a small sip of water with your heart, blood pressure or seizure medication.
- **You must have NOTHING in your mouth 4 hours prior to your procedure.** (This means NO water, ice chips, gum, candy, cough drops, etc.)

**Any contents in your stomach could come up, getting into your lungs while under anesthesia.

GENERAL INSTRUCTIONS

- You may brush your teeth and gargle prior to your test.
- DO NOT wear body LOTION or body JEWELRY.
- Please remove all piercings (tongue, cheek, nose rings, etc) prior to arrival.
- Someone must be with you while you have the procedure and to drive you home. If your driver is **NOT** in our waiting room, your procedure will **NOT** be performed.
- Failure to follow these instructions may cause your procedure to be delayed or cancelled.